

Kevin Bodge, PE PhD

Krista Egan, EI

olsen associates, inc.

and

Mike McGarry

Brevard County

Natural Resources Mgt. Dept.



olsen
associates, inc.
Coastal Engineering


Changes in Beach Profile and Grain Size after Beach Renourishment

February 2018 | Beach Preservation Tech. Conf.




- **How does beach profile shape change after a beach renourishment event?**
- **How does sand grain size change across the profile after a beach renourishment event?**
- **We'll contrast Pre- and Post-Project Conditions.....and first 2 years after a renourishment.**

Brevard County, Florida




Sand Grain Size varies across the Natural Beach


Beach Profile





Sand Grain Size varies across the Natural Beach




Sand Grain Size Distribution









The existing post-project sand matches the pre-project sand by depth zone




The existing post-project sand matches the pre-project sand by depth zone




The existing post-project sand matches the pre-project sand by depth zone




EXPECTED CHANGE IN SAND SIZE ?




Beach Profiles




Beach Profiles




Beach Profiles



Beach Profiles



Beach Profiles




What happens immediately after beach renourishment construction?



We examined:

- Beach profiles: 6 nourished and 3 non-nourished
- Sand grain size at 6 depths along each profile...
...at...
- pre-construction and
- 0-, 3-, 6- 12-, and 24-months post-construction.
 - *54 beach profile surveys, and*
 - *324 sand samples*





Beach Fill Areas —

Nourished (2014) —

Control (No Fill) —

Sand Samples x x x

Nourished Profiles (2013-14)


2013-14 RENOURISHMENT (Brevard County)

Control Profiles (No Fill)

South Reach





olsen
associates, inc.
Coastal Engineering




COMPOSITE PROFILES





COMPOSITE PROFILES





COMPOSITE PROFILES





0 MONTHS

3 MONTHS

6 MONTHS

12 MONTHS

24 MONTHS




NOURISHED


CONTROL




olsen
associates, inc.
Coastal Engineering



BEACH SLOPE ANALYSIS ZONES





SLOPE ANALYSIS: BERM




NR


SR




SLOPE ANALYSIS: INTERTIDAL





NR





SR





SLOPE ANALYSIS: SUBTIDAL




SLOPE ANALYSIS: BERM



SLOPE ANALYSIS: INTERTIDAL






SLOPE ANALYSIS: SUBTIDAL




SURF ZONE WIDTH





SURF ZONE WIDTH




SEABED SAND SAMPLES

R-36

Beach Fill Profile


+3 MSL



GRAIN SIZE DISTRIBUTION




Beach Fill Profile



GRAIN SIZE VS. ELEVATION




Beach Fill Profile





GRAIN SIZE VS. ELEVATION

Control Profile





GRAIN SIZE VS. ELEVATION



MEAN GRAIN SIZE



MEAN GRAIN SIZE



MEAN GRAIN SIZE



Summary – Profiles



- **Post-project beach profile conforms to native pre-project profile, except for steeper beachface slope** (as expected from restoration of eroded beach).
- **Negligible difference in average beach profile shape between nourished and non-nourished, at 0- to 24- months.**
- **Change in nourished beach slopes similar to or lesser than non-nourished profiles from pre- to 12- to 24-months.**
- **Temporal changes in profile slope and surf zone width similar between nourished and non-nourished profiles.**
- **Surf zone widths increasingly converged at 12- to 24-months.**

Summary - Sand



- Existing post-project sand matches native sand, by depth.
- Grain size at nourished berm slightly increased at 0-6 months, returned to pre-renourishment within 12-24 months.
- Little net change in grain size at/below MHW.....changes were less than non-nourished profiles.
- Negligible changes below -9 ft MSL anywhere.
- Temporal variations in grain size are significant -- mostly greater at non-nourished profiles, and obscure trends.....
though there is limited indication of increasing coherence at 12- and 24-months post-construction.

Kevin Bodge, PE PhD

Krista Egan, EI

olsen associates, inc.

and

Mike McGarry

Brevard County

Natural Resources Mgt. Dept.



olsen
associates, inc.
Coastal Engineering

Changes in Beach Profile and Grain Size after Beach Renourishment

February 2018 | Beach Preservation Tech. Conf.